

**PRODUCTION OF PROTEIN HYDROLYSATES
FROM SOY OKARA BY *ASPERGILLUS ORYZAE***

WAN NURNADIA DIYANA BINTI SULAIMAN

UNIVERSITI SAINS MALAYSIA

JUNE 2020



**PUSAT PENGAJIAN
TEKNOLOGI INDUSTRI
UNIVERSITI SAINS
MALAYSIA**

**BORANG PENYERAHAN
DISERTASI MUTAKHIR
SATU (1) NASKAH**

Nama penyelia: PROF. DR. ROSMA BINTI AHMAD

Bahagian: TEKNOLOGI BIOPROSES

Saya telah menyemak semua pembetulan/pindaan yang dilaksanakan oleh
~~Eneik/Puan~~/Cik WAN NURNADIA DIYANA BINTI SULAIMAN

Mengenai disertasinya sebagaimana yang dipersetujui oleh Panel Pemeriksaan di
Viva Vocenya.

2. Saya ingin mengesahkan bahawa saya berpuas hati dengan
pembetulan/pindaan yang dilaksanakan oleh calon.

Sekian, terima kasih.


Tandatangan Signature
Dr. ROSMA AHMAD (PhD)
Associate Professor
Bioprocess Technology Division
School of Industrial Technology
Universiti Sains Malaysia
11800 Minden, Penang

16/07/2020

Tarikh



PRODUCTION OF PROTEIN HYDROLYSATES FROM SOY OKARA BY *ASPERGILLUS ORYZAE*

by

WAN NURNADIA DIYANA BINTI SULAIMAN

A dissertation submitted in the partial fulfilment of the requirements for the degree
of Bachelor of Technology (B. Tech) in the field of Bioprocess Technology
School of Industrial Technology
Universiti Sains Malaysia

June 2020

DECLARATION BY AUTHOR

The dissertation is composed of my original work, and contains no material previously published or written by another person except where due reference has been made in the text. The content of my dissertation is the result of work I have carried out since the commencement of my research project and does not include a substantial part of work that has been submitted to qualify for the award of any other degree or diploma in any university or other tertiary institution.



WAN NURNADIA DIYANA BINTI SULAIMAN

JUNE 2020

ACKNOWLEDGEMENTS

First and foremost, I would like to take this opportunity to express utmost gratitude and special appreciations to my supervisor Prof. Dr. Rosma binti Ahmad for her encouragement and full support throughout this final year project. Her selfless spirit in teaching me step by step helped me a lot in completing this thesis.

Besides, special thanks to Bioprocess laboratory assistant, Mr. Azmaizan bin Yaakub and Mrs. Najmah binti Hamid for assisting me in preparing all those chemicals and apparatus that I need during conducting the project.

Other than that, I would like to take this opportunity to thank all lecturers for their helpful advices. I also want to express my appreciation to my colleagues especially Ummi Syakina binti Sulaiman for helping me along the difficult time.

Furthermore, thank you to my mother Wan Noriza binti Wan Abdullah and my siblings, Wan Nurhusna Auni, Wan Nur'aliala 'Aqilah, and Wan Nurainun Najwa for their prayers, love, financial, and never ending support throughout this project. I am thankful to be with all of you in my life.

Finally, for the rest that I did not mention, believe me you are not forgotten. Thank you from the bottom of my heart.

WAN NURNADIA DIYANA BINTI SULAIMAN

JUNE 2020

TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS	v
TABLE OF CONTENTS	vi
LIST OF TABLES	ix
LIST OF FIGURES	x
LISTS OF SYMBOLS AND ABBREVIATIONS	xi
ABSTRAK	xii
ABSTRACT	xiii
CHAPTER 1 INTRODUCTION	
1.1 Research Background	1
1.2 Problem Statement	3
1.3 Research Objectives	5
CHAPTER 2 LITERATURE REVIEW	
2.1 Soy Bean	6
2.1.1 Soy Okara	7
2.1.2 Chemical Composition of Okara	7
2.1.3 Utilisation of Okara	8
2.2 Solid State Fermentation (SSF)	8
2.2.1 Factors Affecting SSF	9
2.2.1a Moisture Content of Substrate	9
2.2.1b Thickness of Substrate	10
2.3 <i>Aspergillus oryzae</i>	10
2.4 Protein Hydrolysates	11

CHAPTER 3 MATERIALS AND METHODS

3.1 Source of Soy Okara	13
3.2 Proximate Analysis	13
3.2.1 Determination of Moisture Content	13
3.2.2 Determination of Crude Protein Content	14
3.2.3 Determination of Crude Fat Content	16
3.2.4 Determination of Ash Content	17
3.2.5 Determination of Carbohydrate Content	18
3.3 Inoculum Preparation	19
3.3.1 Reactivation of Fungal Stock on Potato Dextrose Agar (PDA)	19
3.3.2 Fungal Reactivation on Soybean	20
3.4 Solid State Fermentation (SSF) Process	21
3.5 Analysis	22
3.5.1 Determination of Amino Acid Concentration	22
3.5.2 Determination of Fungal Growth	23
3.6 Statistical Analysis	24

CHAPTER 4 RESULTS AND DISCUSSION

4.1 Proximate Analysis of Fresh Okara	26
4.1.1 Determination of Moisture Content	26
4.1.2 Determination of Crude Protein Content by using Kjeldahl Method	27
4.1.3 Determination of Crude Fat Content by using Soxhlet Method	29
4.1.4 Determination of Ash Content	30
4.1.5 Determination of Carbohydrate Content (by different)	31

4.2 Growth Profile of <i>Aspergillus oryzae</i> during Okara Fermentation	33
4.2.1 Effect of Okara Thickness on Glucosamine Content during Okara Fermentation with 2.5% Inoculum	34
4.2.2 Effect of Okara Thickness on Glucosamine Content during Okara Fermentation with 5.0% Inoculum	37
4.3 Amino Acid Content	39
4.3.1 Effect of Okara Thickness on Amino Acid Content during Okara Fermentation with 2.5% Inoculum	40
4.3.2 Effect of Okara Thickness on Amino Acid Content during Okara Fermentation with 5.0% Inoculum	43
4.4 Moisture Content during Fermentation	45
4.4.1 Moisture Content of 2.5% Inoculum with Different Thickness	46
4.4.2 Moisture Content of 5.0% Inoculum with Different Thickness	48
CHAPTER 5 CONCLUSIONS AND RECOMMENDATIONS FOR FUTURE RESEARCH	
5.1 Conclusions	50
5.2 Recommendations for Future Research	50
REFERENCES	51
APPENDICES	57

LIST OF TABLES

Table Caption	Page
4.1 Percentage of moisture content	26
4.2 Percentage of nitrogen and protein content	27
4.3 Percentage of crude fat content	29
4.4 Percentage of ash content	30
4.5 Percentage of proximate analysis composition	31
4.6 Increased in glucosamine content during okara fermentation with 2.5% inoculum sample as affected by okara thickness	36
4.7 Increased in glucosamine content during okara fermentation with 5.0% inoculum sample as affected by okara thickness	38
4.8 Increased in amino acid content during okara fermentation with 2.5% inoculum sample as affected by okara thickness	42
4.9 Increased in amino acid content during okara fermentation with 5.0% inoculum sample as affected by okara thickness	44
4.10 Moisture content in 2.5% inoculum sample with 1 cm thickness	46
4.11 Moisture content in 2.5% inoculum sample with 3 cm thickness	47
4.12 Moisture content in 5.0% inoculum sample with 1 cm thickness	48
4.13 Moisture content in 5.0% inoculum sample with 3 cm thickness	49

LIST OF FIGURES

Figure Caption	Page
2.1 The structure of partial hydrolysis	11
3.1 <i>A. oryzae</i> on PDA	20
3.2 White mycelium on agar plate	20
3.3 Before incubation	21
3.4 After incubation (Inoculum for SSF process)	21
3.5 Fermented soybean (inoculum) was cut into pieces for mixing process	22
3.6 The flowchart for research project	25
4.1 Glucosamine standard curve	34
4.2 Time course profile of fungus growth in different thickness of substrate with 2.5% inoculum	35
4.3 Time course profile of fungus growth in different thickness of substrate with 5.0% inoculum	37
4.4 Amino acid (L-leucine) standard curve	40
4.5 Time course profile of amino acid content in different thickness of substrate in 2.5% inoculum	41
4.6 Time course profile of amino acid content in different thickness of substrate in 5.0% inoculum	43

LIST OF SYMBOLS AND ABBREVIATIONS

Symbol	Caption
+	Plus
-	Minus
±	Plus-Minus
%	Percentage
°C	Degree Celcius
&	And
=	Equal
>	More than
Abbreviation	Caption
ml	Millilitre
mg	Milligram
g	Gram
kg	Kilogram
OD	Optical Density
PDA	Potato Dextrose Agar
SmF	Submerged Fermentation
SSF	Solid State Fermentation
USA	United States of America
w/v	Weight Per Volume
v/v	Volume Per Volume
SPSS	Statistical Package for the Social Sciences

PENGHASILAN PROTEIN HIDROLISAT OLEH *ASPERGILLUS ORYZAE*

DARI OKARA KACANG SOYA

ABSTRAK

Soya okara adalah sisa pulpa kacang soya atau hampas kacang soya yang diperoleh setelah menjalani proses pecahan air yang boleh digunakan untuk menghasilkan tauhu atau susu soya. Dalam industri soya, jumlah okara soya yang dihasilkan setiap hari lebih tinggi daripada yang dijangkakan yang mempunyai kandungan protein yang tinggi. Ia dapat meningkatkan pencemaran alam sekitar yang sangat rentan terhadap pembusukan pada okara segar. Hidrolisis protein okara disediakan dengan pembiakkan *Aspergillus oryzae* untuk menghasilkan protein berfungsi untuk makanan dan makanan haiwan. Dalam penyelidikan ini, okara akan dibiakkan dengan peratusan inokulum *A. oryzae* yang berbeza pada inkubasi 25°C selama 72 jam. Pengambilan sampel secara berkala setiap enam jam telah dilakukan dan sampel dianalisis untuk pertumbuhan fungi dengan kaedah glukosamin dan kepekatan asid amino menggunakan ujian ninhidrin. Kandungan kelembapan okara segar adalah 84.51%. Analisis anggaran okara secara kering adalah 33.09% (kandungan protein), 10.66% (kandungan lemak), 0.60% (kandungan abu), dan 55.65% (kandungan karbohidrat). Secara statistik, dengan menggunakan Uji Sampel Bebas, tidak ada perbezaan yang signifikan dengan nilai p 0.280 (2.5%) dan 0.147 (5.0%) untuk kandungan glukosamin sedangkan 0.684 (2.5%) dan 0.344 (5.0%) untuk kandungan asid amino dalam kesan okara untuk kedua-dua ketebalan 1 cm dan 3 cm. Fermentasi okara tidak dipengaruhi oleh ketebalan substrat (1 cm dan 3 cm) dengan 2.5% dan 5.0% inokulum. Kandungan glukosamin maksimum ialah 0.164 mg pada 2.5% dan 0.176 mg pada inokulum 5.0%. Kandungan asid amino tertinggi pada inokulum 2.5% ialah 84.6 μg dan 94.05 μg pada inokulum 5.0%.

PRODUCTION OF PROTEIN HYDROLYSATES FROM SOY OKARA BY *ASPERGILLUS ORYZAE*

ABSTRACT

Soy okara is the soybean pulp or soybean curd residues that obtain after undergoing water-extractable fraction process used to produce bean curd or soymilk. In soy industry, the amount of soy okara produced every day is higher than expected which in high protein content. It can increase the environmental pollutions cause of highly susceptible to putrefaction in fresh okara. Okara protein hydrolysate is prepare by cultivation with *Aspergillus oryzae* to produce functional protein for foods and animal feeds. In this research, okara will be cultivated with different percentage of *A. oryzae* inoculum at 25°C incubation for 72 hours. Periodical sampling at every six hours has been carried out and samples was analysed for fungal growth by glucosamine method and amino acid concentration using ninhydrin test. Moisture content of fresh okara was 84.51%. Proximate analysis in dry basis okara was 33.09% (protein content), 10.66% (fat content), 0.60% (ash content), and 55.65% (carbohydrate content). Statistically, by using Independent Sample T-Test, there was no significant difference with the p-value 0.280 (2.5%) and 0.147 (5.0%) for glucosamine content whereas 0.684 (2.5%) and 0.344 (5.0%) for amino acid content in the effect of okara for both thickness of 1 cm and 3 cm. Okara fermentation was not affected by substrate thickness (1 cm and 3 cm) with 2.5% and 5.0% inoculum. The maximum glucosamine content was 0.164 mg at 2.5% and 0.176 mg at 5.0% inoculum. The highest amino acid content at 2.5% inoculum was 84.6 µg and 94.05 µg at 5.0% inoculum.